

The Araucaria Project. The Distance to the Local Group Galaxy NGC 6822 from Cepheid Variables discovered in a Wide-Field Imaging Survey ¹

Grzegorz Pietrzyński

Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile

Warsaw University Observatory, Al. Ujazdowskie 4, 00-478, Warsaw, Poland

Wolfgang Gieren

Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile

Andrzej Udalski

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Fabio Bresolin

Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA

Rolf-Peter Kudritzki

Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA

Igor Soszyński

Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Michał Szymański

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Marcin Kubiak

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

ABSTRACT

¹Based on observations obtained with the 1.3 m telescope at the Las Campanas Observatory.

We have obtained mosaic images of NGC 6822 in V and I bands on 77 nights. From these data, we have conducted an extensive search for Cepheid variables over the entire field of the galaxy, and we have found 116 such variables with periods ranging from 1.7 to 124 days. We used the long-period (> 5.6 days) Cepheids to establish the period-luminosity relations in V, I and in the reddening-independent Wesenheit index, which are all very tightly defined. Fitting the OGLE LMC slopes in the various bands to our data, we have derived distance values for NGC 6822 in V, I and W_I which agree very well among themselves. Our adopted best distance value from the reddening-free Wesenheit index is 23.34 ± 0.04 (statistical) ± 0.05 (systematic) mag. This value agrees within the combined 1 sigma uncertainties with a previous distance value derived for NGC 6822 by McAlary et al. from near-IR photometry of 9 Cepheids, but our new value is significantly more accurate. We compare the slopes of the Cepheid PL relation in V and I as determined in the five best-observed nearby galaxies, which span a metallicity range from -1.0 to -0.3 dex, and find the data consistent with no metallicity dependence of the PL relation slope in this range. Comparing the magnitudes of 10-day Cepheids with the I-band magnitudes of the TRGB in the same set of galaxies, there is no evidence either for a significant variation of the period-luminosity zero points in V and I. The available data limit such a zero point variation to less than 0.03 mag, in the considered low-metallicity regime.

Subject headings: distance scale - galaxies: distances and redshifts - galaxies: individual: NGC 6822 - galaxies: stellar content - stars: Cepheids

1. Introduction

Nearby galaxies offer a unique laboratory to study in much detail how stellar distance indicators depend on environmental properties like age and metallicity. Calibrating these dependences accurately for the most important stellar candles is a fundamental requirement in order to improve the distance scale. The enormous spread in the distance to the LMC derived from different stellar objects and techniques (Benedict et al. 2002) illustrates the magnitude of the problem, showing that environmental dependences of stellar candles like Cepheids, red clump giants and RR Lyrae stars are not yet under control to the point that the systematic uncertainties in these methods are beat down below a few percent. This affects directly the calibration of the distance scale, and of cosmological parameters linked to it one way or the other. In our ongoing Araucaria Project (see Gieren et al. 2001;

also <http://ifa.hawaii.edu/~bresolin/Araucaria/index.html>), we are addressing the problem by observing large samples of different stellar distance indicators in a sample of nearby galaxies in the Local and Sculptor Groups which cover a broad range in their environmental parameters. Systematic comparison of distance results obtained from the different candles should allow us to recognize and calibrate the dependences on metallicity and age, and possibly other environmental parameters, of the methods of distance determination we are scrutinizing, and to do this with a significantly higher level of accuracy than what has been achieved by other researchers in the literature. First results, for instance on red clump stars, have already been obtained and discussed (Pietrzynski & Gieren 2002; Pietrzynski et al. 2003).

A first step in the exploration of distance indicators in the target galaxies of the Araucaria Project consists in identifying sizeable samples of these objects in each of them. With the advent of the new wide-field mosaic optical detectors, this can be done much more efficiently than in those times when single CCD detectors, or even photographic surveys had to be used to acquire the necessary data, which were then mostly restricted to small parts of these galaxies. For all of our target galaxies it is now possible to obtain high quality CCD photometry for the whole stellar content in a reasonably short time, and therefore significantly improve on results of previous investigations which were derived from the study of small parts of these galaxies only. Our first extensive wide-field survey was performed in the Sculptor spiral galaxy NGC 300, and resulted in the discovery of 117 Cepheids and 12 Cepheid candidates (Pietrzynski et al. 2002) which were recently used to determine the distance to this galaxy from its population of long-period Cepheids (Gieren et al. 2004). We also discovered a considerable number of spectroscopically confirmed blue supergiants in this galaxy from our wide-field survey (Bresolin et al. 2002), which are very useful objects for distance determination out to several Megaparsecs from the Wind Momentum- Luminosity Relationship (Kudritzki et al. 1999), or from the Flux-Weighted Gravity-Luminosity Relationship (Kudritzki et al. 2003; Urbaneja et al. 2004) they have been demonstrated to obey from our NGC 300 data.

As the next target in our survey for distance indicators, we selected the Local Group irregular galaxy NGC 6822. This galaxy is a key object in the Araucaria Project since it contains all the distance indicator classes we study in large numbers, and since it is one of the nearest, and therefore best-resolved galaxies in our target list. Furthermore, the stellar objects are obviously relatively bright, as compared to the more distant galaxies in our target list, facilitating accurate photometry and spectroscopy. NGC 6822 was already subjected to a search for variable stars a long time ago using photographic plates (Hubble 1925, Kayser 1967), which led to discovery of 21 Cepheids. Subsamples of these Cepheids were later observed using CCD detectors in both, the near-IR H band (McAlary et al. 1983

- 9 Cepheids) and optical BVRI bands (Gallart et al. 1996 - 6 Cepheids) in order to improve the distance determination to NGC 6822. Recently, Antonello et al. (2002) monitored a small field of 3.8×3.8 arcmin, located at the centre of NGC 6822, in white light (no filter used), finding 15 new population I Cepheids in this region.

Encouraged by this recent result, we performed a wide-field mosaic imaging survey in NGC 6822 which covers the entire galaxy, with the main goal to discover a significant number of additional Cepheid variables, particularly of long-period Cepheid variables, and other objects we will be studying in our project in forthcoming papers. Using an improved Cepheid database (both in quantity and in quality), our goal was to significantly improve previous distance determinations to this galaxy using the reddening-insensitive Wesenheit magnitude-period relation for Cepheids which we successfully applied in our recent work on NGC 300 (Gieren et al. 2004). In this paper, which is focused on a distance determination from Cepheid variables, we only present the Cepheids we discovered in our mosaic survey. Other types of variables we have also discovered will be presented and discussed in a forthcoming paper, as will be the abundant blue supergiant population in NGC 6822 we have discovered from our data.

In section 2, we present and discuss the observations, calibrations and data reduction procedures. In section 3, we present the catalog of Cepheid variables in NGC 6822 obtained from our data, and in section 4 we use these data to determine the distance to NGC 6822 from the Cepheid PL relation. In section 5, we discuss our result, and its uncertainty, and in section 6 we summarize the main conclusions of this paper.

2. Observations, Reductions and Calibrations

All the data presented in this paper were collected with the Warsaw 1.3-m telescope at Las Campanas Observatory. The telescope was equipped with a mosaic $8k \times 8k$ detector, with a field of view of about 35×35 arcmin and a scale of about 0.25 arcsec/pix. For more instrumental details on this camera, the reader is referred to the OGLE website: <http://ogle.astrouw.edu.pl/~ogle/index.html>. V and I band images of NGC 6822 were secured on 77 different nights. The exposure time was set to 600 seconds in both filters. Most of the observations were obtained in 2002 between July 3 and November 21. Additional observations were made during three photometric nights in May 2003 in order to accurately calibrate the data and improve periods of the detected variable stars.

Preliminary reductions (i.e. debiasing and flatfielding) were done with the IRAF² package. Then, the PSF photometry was obtained for all stars in the same manner as in Pietrzyński, Gieren and Udalski (2002). Independently, the data were reduced with the OGLE III pipeline based on the image subtraction technique (Udalski 2003; Woźniak 2000).

In order to calibrate our photometry onto the standard system our target was monitored during three photometric nights (2003 May 11, 17 and 28) together with some 25 standards from the Landolt fields spanning a wide range of colors ($-0.14 < V-I < 1.43$), and observed at very different airmasses. Since in principle the transformation equations for each of the eight chips may have different color coefficients and zero points, the selected sample of standard stars was observed on each of the individual chips, and transformation coefficients were derived independently for each chip, on each night. Comparison of the photometry obtained on the different nights (see Fig. 1) revealed that the internal accuracy of zero points in both V and I bands is better than 0.01 mag.

To check in detail the possible variation of the zero points in V and I over the mosaic, two fields in the SMC, which contain a huge number of uniformly distributed stars with very accurate photometry established by the OGLE team (Udalski et al. 1999a) were monitored on May 28 together with our target and standard fields, and were calibrated in the same manner as the NGC 6822 observations. Typically, in the selected SMC fields about 200.000 stars were found in common with the OGLE II photometric maps on each of the eight mosaic chips. Comparison of OGLE II and our photometry (see Fig. 2) shows that on average the difference in the zero points between these two photometric sets is smaller than 0.01 mag on each chip. The scatter, being larger than expected from the accuracy of both data sources indicates that there is a small variation of the zero point (up to about 0.05 mag) over the mosaic field, in both filters. In order to account for this effect each chip was divided into $64 \times 512 \times 512$ pixel (about 2×2 arcmin) boxes, and the difference between our photometry and OGLE II data was calculated for each box in both V and I bands. Adopting such a resolution we assure very good statistics in each box, and therefore an accurate (rms smaller than 0.02 mag) determination of our "correction maps". After applying these correction maps in V and I to the SMC fields, the scatter of the difference between the OGLE II and our new mosaic magnitudes for the stars in these fields was reduced down to 0.02-0.03 mag (see Fig. 3), showing that our adopted correction process is indeed improving the accuracy of the photometry derived from the mosaic images. Finally, the NGC 6822 photometry was corrected with these same maps assuring that photometric zero point changes over the mosaic field are kept below 0.03 mag in both, V and I filters.

²IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the NSF.

There are two sources of photometric data available in the literature which allowed us to perform checks on our NGC 6822 photometry. Fig. 4 presents a comparison of the V band photometry of stars from one of the chips of our mosaic camera with photometry published by Bianchi et al. (2001). As one can see, there is no difference between the zero points of these two data sets. Unfortunately, Bianchi et al. (2001) did not provide I band data. Fig. 5 shows the difference between the photometry presented by Gallart et al. (1996) and our results for one of the mosaic chips, in both V and I. In the I band, the difference in the zero points amounts to about 0.03 mag. In the V band, there is practically no offset (e.g. the difference is smaller than 0.005 mag) for stars fainter than 19 mag; however, for brighter stars there is an offset of about 0.03 mag (see upper panel of Fig. 5). Since no such effect is present in the comparison of our data with the photometry obtained by Bianchi et al., and 19 mag in V just corresponds to the saturation limit of long exposures secured by Gallart et al. (1996), we suspect that this jump is associated with a difference between the zero points of the photometry obtained from the long and short exposures of Gallart et al. (1996), which were merged into one database. If we repeat the comparisons using data from other chips of our mosaic camera, the results look very similar.

Summarizing the previous discussion, we may conclude that all the external and internal checks we made indicate that the mean zero points of our data are accurate to 0.03 mag in both V and I, and that their possible variations over the mosaic fields are smaller than 0.03 mag.

3. Cepheid Catalog

All stars observed in NGC 6822 were searched for photometric variations with periods between 0.2 and 100 days, which corresponds to the time baseline covered by our observations, using the AoV algorithm (Schwarzenberg-Czerny 1989). The accuracy of the derived periods is about $1 \times 10^{-4} \times P$. In order to distinguish Cepheids from other types of variable stars several criteria, similar to those used in Pietrzyński et al. (2002), were adopted. All light curves showing the typical shape of Cepheid variables, were approximated by Fourier series of order 4. Then we rejected all candidates with amplitudes in the V band smaller than 0.4 mag, which is approximately the lower limit amplitude for normal classical Cepheids. For the stars remaining in our database, mean V and I band magnitudes were derived by integrating their light curves previously converted onto an intensity scale, and converting the results back to the magnitude scale. Due to the very good quality and phase coverage of our light curves, the statistical accuracy of the derived mean V and I magnitudes was typically better than 0.01 mag. In a final step of selecting the Cepheid

sample in NGC 6822, we deleted objects having mean magnitudes which are widely different (by a full magnitude or more) from those of Cepheids of similar periods, clearly standing off the PL relations presented in section 4 (these and other variables will be studied in a forthcoming paper).

Finally, we ended up with a catalog of 116 Cepheids. Table 1 presents their description: identification, equatorial coordinates, period, time of maximum light in V, mean V and I band magnitudes, their Wesenheit index defined as $W_I = I - 1.55 \times (< V > - < I >)$, and a cross-identification.

Our list of Cepheids contains 9 out of the 11 Cepheids catalogued by Hubble (1925). The stars designed as V10 and V11 by Hubble turned out not to be Cepheids. We also rediscovered all additional Cepheids discovered by Kayser (1967), with the exception of her variable V30. This object is located in a very dense region in NGC 6822 and it was impossible to cross-identify it unambiguously with stars on our images. Six of the Cepheids reported by Antonello et al. (2002) were located between the chips of our mosaic and therefore did not enter our catalog. It should be stressed here that the gaps between the chips of the mosaic installed on the 1.3 m Warsaw telescope are very small (of order of 10 arcsec), so very few Cepheids are expected to be missed due to the gaps. We found that the star designated as V0771 and classified as a W Vir variable with a period of 16.74 days by Antonello et al. seems to be another classical Cepheid. Our data yield a period of 4.6046 days for this star (cep060). Since its location on P-L diagrams fits very well with the locations of other Cepheids with similar periods, we decided to include this star into the present catalog.

Table 2 contains the individual new V and I observations for all the Cepheids in Table 1. The full Table 2 is available in electronic form. In Fig. 6, we are displaying the light curves of several of the Cepheids in our catalog, which are typical in quality for the light curves of other variables of similar periods.

4. PL relations and distance determination

For the purpose of studying the period-luminosity relations defined by the Cepheids in NGC 6822 in V, I and W_I , we excluded all stars having periods larger than 100 days (1 object), and smaller than 5.6 days from the data given in Table 1. A period of 100 days is usually assumed as the upper limit of validity of the PL relation for Cepheids (for a closer discussion of this, see Gieren et al. 2004). The PL relation for Cepheids with periods larger than 100 days becomes nearly flat, which can probably be explained by the nonlinear

pulsation cycle characteristics (Aikawa and Antonello 2000). The short-period cutoff period of 5.6 days was adopted to avoid contamination of our sample by first overtone pulsators. As can be seen from the very impressive PL relations derived from 1333 Cepheids in the LMC, presented by the OGLE team (Udalski et al. 1999c, see their Figure 2), first overtone Cepheids start to appear on the PL diagram shortward of $\log P = 0.75$, which corresponds to our adopted lower limit period of 5.6 days. In the adopted period range from 5.6 to 100 days, three additional stars, cep026, cep029 and cep057, were rejected from our sample because of their strongly deviating location on the P-L diagrams. These 3 objects have mean Wesenheit magnitudes more than 3 sigma different from the mean magnitudes of Cepheids with similar periods. Our final sample of Cepheids used for the analysis of the PL relations in NGC 6822 contains 42 objects. As a consequence of our selection process, there is a very high probability that all of these variables are classical fundamental mode pulsators. We also note that our final Cepheid sample contains a very considerable number of long-period variables (21 with periods longer than 10 days), which are crucial for an accurate distance determination with the period-luminosity relation.

Fits to a straight line to our data yield the following slopes for the PL relations: -2.66 ± 0.07 , -2.92 ± 0.05 and -3.32 ± 0.04 in V, I and W_I , respectively. These values agree very well with the corresponding OGLE LMC slopes of -2.775 , -2.977 and -3.300 (Udalski 2000). In particular there is excellent agreement for the slope of the reddening-free W_I index, crucial for the distance determination. It is worthwhile noticing, at this point, that in recent work basically identical slopes for this relation were obtained for several galaxies which span a large range in metallicities (e.g. SMC - Udalski et al. 1999b, IC 1613 - Udalski et al. 2001, NGC 300 - Gieren et al. 2004), indicating that the Wesenheit (V-I) PL relation must be substantially metallicity independent, making it a very powerful tool for distance determination. A full analysis of the constancy of the PL slopes in different photometric bands will be undertaken once we have obtained PL relations for all the galaxies of the Araucaria Project.

Given the good agreement of the PL slopes in the different bands derived for NGC 6822 from our data, with the LMC OGLE slopes, we chose to adopt the extremely well determined OGLE slopes to derive the distance to NGC 6822. This leads to the following equations:

$$V = -2.775 \log P + (23.024 \pm 0.033)$$

$$I = -2.977 \log P + (22.118 \pm 0.026)$$

$$W_I = -3.300 \log P + (20.706 \pm 0.021)$$

Adopting 18.50 mag as the distance modulus to the LMC, $E(B-V) = 0.36$ mag as the sum of foreground and intrinsic reddenings for NGC 6822 (McAlary et al. 1986), and the reddening law of Schlegel et al. (1998) ($A_V = 3.24 E(B-V)$, $A_V = 1.96 E(B-V)$) we obtain the following distance moduli for NGC 6822 in the three different bands we study:

$$(m - M)_0 (W_I) = 23.338 \text{ mag}$$

$$(m - M)_0 (I) = 23.319 \text{ mag}$$

$$(m - M)_0 (V) = 23.292 \text{ mag}$$

As can be seen, the results from the three different bands are in excellent agreement. Possible sources of uncertainty affecting these values are discussed in the next section. The very slightly shorter distances obtained in the V and I bands may be caused by a slight overestimation of the adopted reddening. Indeed, reducing the assumed reddening by just 0.01 mag we would get practically the same distances (within 0.01 mag) from all three bands.

5. Discussion

Before arriving at our best value for the distance to NGC 6822, together with the associated statistical and systematical errors (see the next section), we will now discuss how the potential dependence of the PL relation on metallicity, blending with unresolved companion stars, and the uncertainty of the assumed reddening may affect our result. Another contribution to the total error budget, the uncertainty of the zero point of our photometry, was already discussed in detail in section 2. We will not discuss here the probably largest contribution to the systematic error of the distance to NGC 6822, which is the current uncertainty of our adopted LMC distance modulus. The reader is referred to the paper of Benedict et al. (2002) for a modern review of distance determinations to the LMC. In our paper, we assume a true distance modulus to the LMC of 18.50 mag in order to place our distance of NGC 6822 onto the same scale as the Cepheid distances to some 30 galaxies derived by the HST Key Project for the Extragalactic Distance Scale (Freedman

et al. 2001), and to be consistent with our recent distance determination from Cepheid variables for NGC 300 (Gieren et al. 2004).

5.1. Dependence of the PL relation on metallicity

In order to investigate the possible dependence of the Cepheid period-luminosity relation on the metallicity of the Cepheids, from a purely empirical point of view, we can compare the well-determined period-luminosity relation in NGC 6822 of this paper to the PL results obtained for other galaxies of different metallicities. According to the high spectral resolution study of Venn et al. (2001), the mean metallicity of the young stellar population in NGC 6822 is $[\text{Fe}/\text{H}] = -0.5$ dex, intermediate between LMC and SMC. We select for comparison only those nearby galaxies for which large samples of Cepheids have very well-measured light curves in the V and I bands, which have allowed to establish the coefficients of their respective PL relations with high accuracy. The list of these galaxies currently includes the LMC, SMC, IC 1613 and NGC 300 (and will be expanded in the course of our project). It is worthwhile noticing that it is reasonable to assume (see the next section) that for all these well-resolved galaxies blending does not significantly affect the measured Cepheid brightnesses, and therefore the slopes and zero points which were obtained from the analysis of the respective PL relations.

As we already mentioned in the previous section, the slopes of the PL relation in the V, I and W_I bands in NGC 6822 agree very well with the values of the slopes derived from the Cepheids in the LMC (Udalski et al. 1999b, Udalski 2000), SMC (Udalski et al. 1999b, Udalski 2000), IC 1613 (Udalski et al. 2001) and NGC 300 (Gieren et al. 2004). The differences of the observed slopes in these five galaxies, in any of these bands, do never deviate by more than about 1.5 sigma of the respective determinations, which we interpret as a clear sign that from the excellent Cepheid data available for this well-observed set of galaxies, no significant difference in the slope of the PL relation can be established. Recalling that these galaxies have mean metallicities of their Cepheid populations in the range $-1.0 < [\text{Fe}/\text{H}] < -0.3$, our current work on NGC 6822 confirms and strengthens the earlier finding of Udalski et al. (2001) that the slope of the Cepheid PL relation in the V and I bands seems to be universal in this metallicity range. As a note of caution, however, we note that the situation may change for the higher-metallicity regime, where careful studies of Galactic Cepheids of near-solar mean metallicity have provided some evidence for steeper PL slopes (e.g. Storm et al. 2004; Gieren et al. 1998). Studies to confirm this later finding must continue, at an ever increasing level of accuracy.

To verify if and how the absolute zero point of the Cepheid PL relation depends on

the metal content we decided to use the I-band magnitude of the tip of the red giant branch (TRGB) as the reference brightness for Cepheids. There are strong reasons to believe that such an approach leads to very robust results. I-band TRGB magnitudes can be derived very accurately from the excellent number statistics provided by wide-field surveys of nearby galaxies; the TRGB brightness is similar to the Cepheid brightness and, most importantly for our current discussion, the TRGB I-band magnitude does not depend on metallicity for a wide range of metallicities (Ballazzini, Ferraro and Pacino 2001, and references therein). In particular, the comparison of the I-band magnitudes of Cepheids and the TRGB measured from the same data is purely differential and the errors of the photometric zero point and the adopted reddening should cancel out, and therefore such a comparison can be done very precisely.

To determine the I-band magnitude of the TRGB in NGC 6822 from our data we selected the red giants from their positions on the CMD diagram. Fig. 10 presents the histogram of the brightness of the selected red giants, constructed with 0.04 mag bins. The location of the TRGB is clearly depicted in this diagram at 19.83 ± 0.03 mag. This value agrees very well with the TRGB magnitude of 19.8 ± 0.1 mag previously determined for NGC 6822 by Gallart et al. (1996). Finally, and following Udalski et al. (2001), we can derive the difference between the I-band magnitude of the TRGB and the I-band magnitude of Cepheids with a period of 10 days. The value for NGC 6822 from our data is 0.68 ± 0.05 mag. As can be seen from Fig. 11, which compares this difference to the corresponding values observed for the LMC, SMC and IC 1613 (see Udalski et al. (2001)), the difference value is constant to within 0.03 mag for these galaxies, a result which strongly supports the idea that, apart from the slopes in V and I, the absolute I-band zero point of the Cepheid PL relation is universal in the metallicity range from -1.0 and -0.3 dex.

One may therefore conclude that adopting the OGLE LMC slopes and ZPs for the distance determination to NGC 6822 we do not introduce any significant systematic error, given the small metallicity difference between the LMC (-0.3 dex) and NGC 6822 (-0.5 dex).

5.2. Blending

As was shown by Mochejska et al. (2000, 2001), Cepheid magnitudes obtained from ground-based data with a typical resolution of 1.5 arcsec, may be systematically overestimated up to 0.07 mag (for Cepheids with periods longer than 10 days) due to crowding for relatively nearby galaxies like M31 and M33. Since NGC 6822 is located at a shorter distance of roughly a 60 % of the distance of M 33, and our images are of higher resolution than those of M 33 used by Mochejska et al., it is reasonable to expect that the

effect of blending on the Cepheid magnitudes in NGC 6822 reported in this paper will be smaller than 0.07 mag. Indeed, Udalski et al. (2001) comparing ground-based and HST magnitudes of Cepheids in IC 1613, a galaxy of the same type and of similar distance as NGC 6822, concluded that blending does not affect the ground-based measurements by more than 0.04 mag. Another argument in favour of a very small effect of crowding on our ground-based Cepheid magnitudes in NGC 6822 comes from a simulation made by Stanek and Udalski (1999), who, using the OGLE data, analyzed how the magnitudes of the LMC Cepheids observed at a worse resolution (e.g. larger distance) would be affected by crowding. Since the LMC is a galaxy of the same type as NGC 6822, we can use their results to predict how strong a blending effect one should expect for Cepheids in NGC 6822. As can be seen from Fig. 4 in Stanek and Udalski (1999), the increased blending would affect the LMC Cepheid magnitudes at an average level of 0.03 mag, only. In the light of the above discussion it is reasonable to assume that crowding should have an effect on our distance determination presented in this paper which does not exceed 0.04 mag, in the sense that the true distance to NGC 6822 should not be more than a maximum of 0.04 mag larger than our derived result.

5.3. Reddening

Unfortunately, NGC 6822 is located relatively close to the Galactic plane ($b = -18$ deg), which results in a relatively large, and possibly differential reddening across the galaxy. However, the tightness of the PL relations obtained in this paper for the Cepheids in NGC 6822 (see Figs 5-7) clearly suggests that the differential reddening to the Cepheids is not strong. Furthermore, in our approach we have circumvented the problem of the lack of information about the individual reddenings of the Cepheids by using the reddening-free Wesenheit index W_I . It is remarkable that assuming a constant $E(B-V) = 0.36$ mag as advocated by McAlary et al. (1983) we obtain fully consistent distances from the PL relations constructed in the V, I and W_I bands, which again indicates that differential reddening is not an important issue in NGC 6822, and that the adopted mean reddening must be very nearly correct.

Finally, our best distance modulus for NGC 6822 of 23.34 mag agrees quite well with the distance modulus of 23.47 ± 0.11 mag as derived by McAlary et al. (1983) from observations of 9 Cepheids in the near-infrared H-band, where the influence of reddening is significantly reduced as compared to the optical bands used in our study. This supports our conclusion that uncertainties due to reddening have a very small, and probably negligible influence on our distance result.

6. Conclusions

Our best distance value for NGC 6822, which comes from the reddening-free Wesenheit magnitudes is 23.34 ± 0.04 (statistical) ± 0.05 (systematic). The quoted statistical error contains the quadratically combined uncertainties of the OGLE and our fits to the PL relations of the LMC and NGC 6822, respectively, while the systematic uncertainty includes the quadratically combined uncertainties of the zero points of both photometric studies, and a contribution due to blending. As remarked before, our distance value does *not* take into account the systematic uncertainty of the adopted LMC distance of 18.50 mag. Should future research provide an improved determination of the distance to the LMC, our derived distance to NGC 6822 can be easily adjusted to such a change. As extensively discussed in the previous section, our NGC 6822 distance value seems very little affected by any systematic error associated with a possible dependence of the Cepheid PL relation on metallicity, and with reddening. Taking this into account, plus the fact that our new distance determination comes from a large sample of Cepheids (about 6 times larger than the samples of Cepheids used for previous distance determinations to NGC 6822) with very well established light curves and periods, we conclude that the presented result is the to-date most robust distance determination to this galaxy.

A comparison of the slopes of the V- and I-band PL relations obtained from the currently best-observed galaxies (LMC, SMC, IC 1613, NGC 6822 and NGC 300) strongly suggests that the slopes in both bands do not depend on metallicity, in the range $-1.0 < [\text{Fe}/\text{H}] < -0.3$ defined by these galaxies. A comparison of the I-band TRGB magnitudes with the magnitudes of Cepheids at a pulsation period of 10 days in the same sample of galaxies does not provide any evidence for a significant variation of the PL zero points in V and I with metallicity either. This finding evidently strengthens our confidence that Cepheid variables are excellent standard candles, at least when they are used in the low-metallicity regime.

We are grateful to Las Campanas Observatory, and to the CNTAC for providing the large amounts of telescope time which were necessary to complete this project. GP and WG gratefully acknowledge financial support for this work from the Chilean Center for Astrophysics FONDAF 15010003. WG also acknowledges support from the Centrum fuer Internationale Migration und Entwicklung in Frankfurt/Germany who donated the Sun Ultra 60 workstation on which a substantial part of the data reduction and analysis for this project was carried out. Support from the Polish KBN grant No 2P03D02123 and BST grant for Warsaw University Observatory is also acknowledged.

REFERENCES

- Aikawa and Antonello, 2000, *A&A*, 363, 601
- Antonello, E., Fugazza, D., Mantegazza, L., Stefanon, M., Covino, S., 2002, *A&A*, 386, 860
- Bellazzini, M., Ferraro, F.R., and Pacino, E., 2001, *ApJ*, 556, 635
- Benedict, G.F., McArthur, B.E., Fredrick, L.W., et al., 2002, *AJ*, 123, 473
- Bianchi, L., Scuderi, S., Massey, P., Romaniello, M., 2001, *AJ*, 121, 2020
- Bresolin, F., Gieren, W., Kudritzki, R.-P., Pietrzyński, G. and Przybilla, N., 2002, *ApJ*, 567, 277
- Freedman, W.L. et al., 2001, *ApJ*, 553, 47
- Gallart, C., Aparicio, A., and Vilchez, J.M., 1996, *AJ*, 112, 1928
- Gieren, W.P., Fouqué, P., and Gómez, M., 1998, *ApJ*, 496, 17
- Gieren, W., Geisler, D., Richtler, T., Pietrzyński, G., Dirsch, B., 2001, *The Messenger*, 106, 15
- Gieren, W., Pietrzyński, G., Walker, A., Bresolin, F., Minniti, D., Kudritzki, R.P., Udalski, A., Soszyński, I., Fouqué, P., Storm, J., and Bono, G., 2004, *AJ*, in press
- Hubble, E., 1925, *ApJ*, 62, 409
- Kayser, S.E., 1967, *AJ*, 72, 134
- Kudritzki, R.P., Puls, J., Lennon, D.J., Venn, K.A., Reetz, J., Najarro, F., McCarthy, J.K., Herrero, A., 1999, *A&A*, 350, 970
- Kudritzki, R.P., Bresolin, F., Przybilla, N., 2003, *ApJ*, 582, 83
- McAlary, W.Ch., Madore, B., McGonegal, R., McLaren, R.A., and Welch, D.L., 1983, *ApJ*, 273, 539
- Mochejska, B.J., Macri, L.M., Sasselov, D.D., and Stanek, K.Z., 2000, *AJ*, 120, 810
- Mochejska, B.J., Macri, L.M., Sasselov, D.D., and Stanek, K.Z., 2001, *astro-ph/0103440*
- Pietrzyński, G., Gieren, W., Fouqué, P., and Pont, F., 2002, *AJ*, 123, 789
- Pietrzyński, G., and Gieren, W., 2002, *AJ*, 124, 2633
- Pietrzyński, G., Gieren, W., and Udalski, A., 2002, *PASP*, 114, 298
- Pietrzyński, G., Gieren, W., and Udalski, A., 2003, *AJ*, 125, 2494
- Schlegel, D.J., Finkbeiner, D.P., and Davis, M., 1998, *ApJ*, 500, 525
- Schwarzenberg-Czerny, A., 1989, *MNRAS*, 241, 153

- Stanek, K.Z., and Udalski A., 1999, astro-ph/9909346
- Storm, J., Carney, B.W., Gieren, W.P., Fouqué, P., Latham, D.W. and Fry, A.M., 2004, A&A, 415, 531
- Udalski, A., 2000, Acta Astron., 50, 279
- Udalski, A., 2003, Acta Astron., 53, 291
- Udalski, A., Soszyński, I., Szymański, M., Kubiak, M., Pietrzyński, G., Woźniak, P., and Żebruń, K., 1999c, Acta Astron., 49, 45
- Udalski, A., Soszyński, I., Szymański, M., Kubiak, M., Pietrzyński, G., Woźniak, P., and Żebruń, K., 1999a, Acta Astron., 49, 223
- Udalski, A., Szymański, M., Kubiak, M., Pietrzyński, G., Soszyński, I., Woźniak, P., and Żebruń, K., 1999b, Acta Astron., 49, 201
- Udalski, A., Wyrzykowski, Ł, Pietrzyński, G., Szewczyk, O., Szymański, M., Kubiak, M., Soszyński, I., and Żebruń, K., 2001, Acta Astron., 51, 222
- Urbaneja, M.A., Herrero, A., Bresolin, F., Kudritzki, R.-P., Gieren, W., Puls, J., Przybilla, N., Najarro, F. and Pietrzyński, G., 2004, ApJ, submitted
- Venn, K.A., et al. 2001, ApJ, 547, 765
- Woźniak, P., 2000, Acta Astron., 50, 421

Table 1. Cepheids in NGC 6822

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀ -2450000	< V > [mag]	< I > [mag]	< W _I > [mag]	Remarks
cep001	19:45:02.01	-14:47:33.1	123.9	2486.8008	17.864	16.464	14.294	H-V13, A-V1292
cep002	19:44:56.82	-14:53:15.7	65.32	2566.5870	17.677	16.428	14.492	H-V7
cep003	19:44:55.23	-14:47:12.7	37.461	2541.6318	18.688	17.423	15.462	H-V2, A-V3873
cep004	19:44:56.76	-14:51:18.5	34.663	2565.6012	18.865	17.578	15.583	K-V28
cep005	19:45:25.05	-14:58:44.8	32.419	2492.7951	18.840	17.664	15.841	
cep006	19:45:15.81	-14:55:24.2	31.868	2577.5338	18.560	17.397	15.594	K-V29
cep007	19:44:59.72	-14:47:10.8	30.512	2489.7943	19.014	17.736	15.755	H-V1, V3736
cep008	19:44:53.05	-14:46:56.6	29.211	2502.7439	19.219	17.931	15.935	H-V3, A-V2374
cep009	19:45:43.30	-14:45:43.1	21.120	2529.7086	19.416	18.276	16.509	
cep010	19:45:02.22	-14:53:44.2	19.960	2599.5192	19.454	18.249	16.381	H-V6
cep011	19:44:52.42	-14:47:36.4	19.887	2770.8827	19.878	18.523	16.423	A-V1286
cep012	19:45:08.26	-14:53:49.9	19.602	2566.5870	19.759	18.457	16.439	
cep013	19:44:47.18	-14:49:21.9	19.295	2489.7943	19.715	18.583	16.828	K-V17
cep014	19:45:06.44	-14:51:04.8	18.339	2500.7632	19.562	18.352	16.476	K-V21
cep015	19:44:56.56	-14:50:34.3	17.344	2502.7439	19.500	18.371	16.621	H-V4
cep016	19:44:51.32	-14:54:19.1	16.960	2578.5411	19.872	18.699	16.881	
cep017	19:44:53.09	-14:48:39.8	16.855	2564.6135	20.117	18.780	16.708	H-V9, A-V0734
cep018	19:45:02.54	-14:51:06.8	13.872	2548.6332	19.890	18.672	16.784	H-V5
cep019	19:44:50.35	-14:49:13.6	11.164	2511.7407	19.978	18.823	17.033	
cep020	19:45:23.48	-14:46:19.0	10.925	2575.5684	19.525	18.677	17.363	
cep021	19:45:46.63	-14:56:42.4	10.783	2490.7696	19.865	18.899	17.402	
cep022	19:44:50.16	-14:52:29.0	10.277	2578.5411	20.544	19.382	17.581	
cep023	19:45:04.55	-14:55:08.5	9.5731	2492.7951	20.205	19.124	17.448	
cep024	19:44:50.94	-14:48:37.3	9.3664	2574.5653	20.453	19.256	17.401	K-V25, A-V0729
cep025	19:45:00.46	-14:47:04.8	8.9367	2464.8644	20.933	19.236	16.606	A-V3738
cep026	19:45:04.46	-14:52:53.5	8.4670	2518.7165	20.249	19.131	17.398	
cep027	19:45:04.64	-14:53:40.6	8.3912	2497.7874	20.561	19.387	17.567	
cep028	19:45:05.65	-14:53:21.0	7.2085	2554.6080	19.920	18.529	16.373	
cep029	19:45:02.93	-14:51:57.3	7.1054	2471.8358	20.918	19.749	17.937	
cep030	19:44:52.00	-14:45:52.1	6.8842	2521.7141	20.965	19.771	17.920	
cep031	19:44:51.12	-14:47:20.6	6.8665	2489.7943	20.819	19.717	18.009	A-A1433
cep032	19:45:27.86	-14:53:52.4	6.7650	2517.7361	20.727	19.743	18.218	
cep033	19:45:01.63	-14:49:21.8	6.7356	2496.7490	20.959	19.675	17.685	A-V0372

Table 1. Cepheids in NGC 6822 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀ -2450000	< V > [mag]	< I > [mag]	< W _I > [mag]	Remarks
cep034	19:45:08.76	-14:49:10.7	6.5391	2460.8596	20.739	19.541	17.684	
cep035	19:45:01.58	-14:43:10.1	6.2965	2770.8827	20.754	19.665	17.977	
cep036	19:45:19.50	-14:45:58.4	6.1457	2463.8594	20.568	19.625	18.163	
cep037	19:44:51.06	-14:46:06.0	6.1387	2545.6034	21.074	20.030	18.412	
cep038	19:44:30.71	-14:51:22.1	6.0637	2548.6332	20.904	19.817	18.132	
cep039	19:45:23.88	-14:46:33.3	6.0011	2458.8719	20.734	19.691	18.074	
cep040	19:45:36.21	-14:54:56.5	5.9793	2552.6243	21.236	19.424	16.615	
cep041	19:45:04.66	-14:49:39.5	5.9760	2516.7308	20.498	19.543	18.063	A-V0271
cep042	19:44:12.78	-14:41:08.3	5.7100	2464.8644	20.811	19.857	18.378	
cep043	19:45:00.93	-14:47:50.1	5.6450	2463.8594	20.913	19.835	18.164	
cep044	19:44:33.77	-14:39:36.1	5.5905	2557.6145	20.919	19.862	18.224	
cep045	19:44:44.05	-14:51:26.8	5.5621	2463.8594	21.031	19.941	18.251	
cep046	19:44:54.16	-14:49:47.9	5.5518	2787.8786	20.992	19.953	18.343	
cep047	19:45:23.68	-14:49:03.5	5.3611	2584.5470	20.787	19.850	18.398	
cep048	19:44:49.45	-14:46:57.0	5.1764	2591.5357	21.523	20.333	18.488	
cep049	19:45:21.67	-14:53:56.9	5.1701	2548.6332	21.022	20.252	19.059	
cep050	19:45:08.19	-14:41:27.5	5.1285	2498.7581	21.146	20.214	18.769	
cep051	19:45:01.67	-14:51:27.8	5.1090	2579.5495	20.686	19.898	18.677	
cep052	19:45:08.45	-14:46:43.1	5.0970	2488.7265	20.998	20.046	18.570	
cep053	19:44:25.34	-14:43:39.4	5.0773	2518.7165	21.335	20.234	18.527	
cep054	19:45:12.08	-14:44:43.5	5.0560	2546.6257	21.056	20.162	18.776	
cep055	19:45:11.37	-14:49:53.1	4.8955	2496.7490	21.086	20.021	18.370	
cep056	19:44:56.64	-14:45:55.8	4.8822	2529.7086	21.297	20.397	19.002	
cep057	19:45:08.01	-14:54:12.2	4.8305	2562.5883	20.880	19.992	18.616	
cep058	19:44:53.84	-14:49:06.7	4.6860	2486.8008	21.542	20.180	18.069	
cep059	19:44:54.24	-14:43:18.0	4.6345	2521.7141	21.050	20.057	18.518	
cep060	19:44:37.93	-14:48:31.0	4.6046	2520.7108	21.196	20.100	18.401	A-V0771
cep061	19:44:59.35	-14:47:45.3	4.5480	2458.8719	21.275	20.031	18.103	A-V3186
cep062	19:44:59.64	-14:44:34.2	4.5264	2593.5212	21.094	20.053	18.439	
cep063	19:44:50.75	-14:46:20.1	4.4495	2526.6949	21.005	19.701	17.680	
cep064	19:44:54.70	-14:51:10.9	4.3602	2555.6347	21.121	19.973	18.194	
cep065	19:44:19.00	-14:40:48.8	4.2612	2574.5653	21.067	20.066	18.514	
cep066	19:44:10.57	-14:45:32.8	4.2330	2496.7490	21.193	20.270	18.839	
cep067	19:45:08.98	-14:54:13.8	4.2327	2488.7265	21.205	20.147	18.507	

Table 1. Cepheids in NGC 6822 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀ -2450000	< V > [mag]	< I > [mag]	< W _I > [mag]	Remarks
cep068	19:44:51.23	-14:50:47.8	4.2294	2556.6132	21.137	20.214	18.783	
cep069	19:45:05.54	-14:52:20.7	4.2188	2555.6347	21.405	20.310	18.613	
cep070	19:44:50.69	-14:50:02.5	4.1123	2556.6132	21.375	20.364	18.797	
cep071	19:45:49.90	-14:53:41.9	4.0912	2493.7883	21.412	20.359	18.727	
cep072	19:44:55.25	-14:49:22.2	4.0195	2552.6243	21.334	20.459	19.103	A-V0410
cep073	19:44:49.90	-14:51:42.1	3.9879	2556.6132	21.523	20.634	19.256	
cep074	19:45:37.87	-14:54:11.1	3.9362	2487.7294	21.245	20.365	19.001	
cep075	19:44:47.97	-14:44:28.6	3.8970	2517.7361	21.427	20.416	18.849	
cep076	19:44:57.33	-14:49:10.8	3.7880	2490.7696	20.007	19.255	18.089	A-V0444
cep077	19:44:59.98	-14:51:27.7	3.7751	2500.7632	21.119	19.919	18.059	
cep078	19:44:59.37	-14:50:19.1	3.7180	2547.6173	21.072	20.028	18.410	
cep079	19:44:46.87	-14:50:26.6	3.7079	2526.6949	21.080	19.531	17.130	
cep080	19:45:02.95	-14:43:30.9	3.6800	2488.7265	21.822	20.560	18.604	
cep081	19:44:57.24	-14:49:51.4	3.6479	2526.6949	21.635	20.587	18.963	A-V0171
cep082	19:44:36.14	-14:43:35.0	3.6190	2490.7696	21.427	20.494	19.048	
cep083	19:44:49.96	-14:54:29.5	3.5845	2460.8596	21.313	20.059	18.115	
cep084	19:44:09.68	-14:41:23.1	3.5468	2770.8827	21.133	20.431	19.343	
cep085	19:45:15.76	-14:44:08.3	3.5078	2463.8594	21.278	20.402	19.044	
cep086	19:44:53.49	-14:50:21.4	3.4584	2546.6257	21.633	20.533	18.828	
cep087	19:44:51.80	-14:53:00.9	3.3700	2546.6257	21.501	20.834	19.800	
cep088	19:44:11.68	-14:49:37.0	3.3659	2563.5980	21.569	21.790	22.133	
cep089	19:45:14.27	-14:53:14.9	3.3032	2520.7108	21.544	20.449	18.752	
cep090	19:45:47.48	-14:54:24.7	3.2916	2770.8827	21.430	20.504	19.069	
cep091	19:44:36.19	-14:45:07.1	3.2780	2550.6265	21.511	20.501	18.935	
cep092	19:44:02.24	-14:38:02.0	3.2273	2498.7581	21.178	99.999	99.999	
cep093	19:45:12.01	-14:53:36.3	3.2069	2492.7951	21.591	20.712	19.350	
cep094	19:44:58.35	-14:50:31.3	3.1756	2546.6257	21.525	20.413	18.689	
cep095	19:44:45.50	-14:49:29.7	3.1093	2576.5305	21.637	20.824	19.564	
cep096	19:45:09.50	-14:44:21.9	2.8979	2496.7490	21.608	20.597	19.030	
cep097	19:44:57.74	-14:52:01.4	2.8486	2502.7439	21.139	20.226	18.811	
cep098	19:44:43.98	-14:51:53.1	2.6760	2559.5401	21.913	21.141	19.944	
cep099	19:44:58.34	-14:53:18.2	2.6684	2547.6173	21.944	20.797	19.019	
cep100	19:44:49.56	-14:45:10.1	2.5950	2547.6173	21.077	20.229	18.915	
cep101	19:45:06.90	-14:53:21.4	2.5937	2578.5411	21.391	19.744	17.191	

Table 1. Cepheids in NGC 6822 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀ -2450000	< V > [mag]	< I > [mag]	< W _I > [mag]	Remarks
cep102	19:45:21.13	-14:48:33.2	2.5839	2577.5338	21.554	20.343	18.466	
cep103	19:44:54.87	-14:50:26.2	2.5390	2517.7361	21.849	20.481	18.361	
cep104	19:44:34.06	-14:38:21.5	2.4540	2577.5338	21.920	99.999	99.999	
cep105	19:44:21.82	-14:41:49.7	2.4536	2546.6257	21.760	20.987	19.789	
cep106	19:45:51.25	-14:49:09.2	2.3725	2593.5212	19.162	17.719	15.482	
cep107	19:45:08.83	-14:46:10.1	2.3571	2500.7632	21.672	20.724	19.255	
cep108	19:45:41.90	-14:36:17.2	2.3124	2557.6145	21.887	21.163	20.041	
cep109	19:45:10.28	-14:53:02.0	2.1755	2471.8358	21.993	21.372	20.409	
cep110	19:45:51.25	-14:57:43.7	2.0915	2554.6080	22.053	20.838	18.955	
cep111	19:45:35.61	-14:49:11.2	2.0772	2555.6347	21.162	20.410	19.244	
cep112	19:45:39.71	-14:38:32.8	2.0740	2559.5401	22.253	21.395	20.065	
cep113	19:45:09.73	-14:55:09.6	2.0697	2487.7294	20.942	19.860	18.183	
cep114	19:45:33.97	-14:54:53.7	2.0645	2559.5401	21.255	20.518	19.376	
cep115	19:44:38.11	-14:52:41.1	1.8683	2579.5495	22.262	21.432	20.145	
cep116	19:45:05.95	-14:51:23.0	1.7092	2486.8008	21.547	20.171	18.038	

Table 2. Individual V and I Observations

object	filter	HJD-2450000	mag	σ_{mag}
cep001	V	2458.87193	18.044	0.014
cep001	V	2460.85964	18.059	0.018
cep001	V	2463.85945	18.124	0.013
cep001	V	2464.86437	18.143	0.013
cep001	V	2465.85531	18.149	0.014
cep001	V	2468.85143	18.207	0.014
cep001	V	2470.84386	18.237	0.015
cep001	V	2471.83581	18.231	0.014
cep001	V	2482.70757	18.327	0.032
cep001	V	2483.78027	18.358	0.024
cep001	V	2486.80078	18.370	0.023
cep001	V	2487.72942	18.367	0.021
cep001	V	2488.72654	18.352	0.019
cep001	V	2489.79432	18.366	0.012
cep001	V	2490.76962	18.337	0.020
cep001	V	2492.79510	18.313	0.020
cep001	V	2493.78829	18.344	0.016
cep001	V	2495.79027	18.258	0.016
cep001	V	2496.74902	18.248	0.013
cep001	V	2497.78736	18.231	0.015

Note. — The complete version of this table is in the electronic edition of the Journal. The printed edition contains only the the first 20 measurements in V band for the Cepheid variable cep001.




Fig. 1.— Comparison of V and I band magnitudes of stars in NGC 6822 observed and calibrated on two different nights on one of the chips of the mosaic camera. An excellent (always better than 0.01 mag) internal agreement is demonstrated.

Fig. 2.— Comparison of OGLE II, and our new mosaic photometry in the SMC fields (see text), for one chip. Additional scatter (up to 0.05 mag) is clearly seen, and most probably it is caused by variations of the zero points over the mosaic field.

Fig. 3.— Comparison of OGLE II, and our new photometry in the SMC fields, after correcting our data for variations of the zero point over the mosaic field- see text for details. This plot demonstrates that our correction maps are useful to correct the variations of the zero point over the mosaic down to about 0.03 mag.




Fig. 4.— Comparison of our NGC 6822 V-band photometry with the photometric data published by Bianchi et al. (2001).




Fig. 5.— Comparison of our NGC 6822 V- and I-band photometry with the data published by Gallart et al. (1996).




Fig. 6.— Phased V- and I-band light curves for some Cepheids of different periods in our NGC 6822 catalog. These light curves are representative for the light curves of other Cepheid variables of similar periods.




Fig. 7.— The period-luminosity relation for NGC 6822 Cepheids in the V band. Open circles denote variables not used in the distance determination because some of these stars might be overtone pulsators. The slope of the relation was adopted from the LMC Cepheids (OGLE II).




Fig. 8.— Same as Fig. 6, for the I band.




Fig. 9.— Same as Fig. 6, for the reddening-independent (V-I) Wesenheit magnitudes. We note that using all the stars (open circles included) in our distance determination to NGC 6822, the result changes in a negligible way.




Fig. 10.— Luminosity function of the red giant branch in NGC 6822 constructed from our mosaic data using bins of 0.05 mag. The arrow shows the position of the TRGB magnitude.




Fig. 11.— Comparison of the I-band brightness of Cepheids with a period of 10 days, and the I-band magnitude of the tip of the red giant branch for four galaxies covering a wide range in metallicities, and having excellent data for the determination of these quantities.

This figure "pietrzynski.fig2a.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0408572v1>

This figure "pietrzynski.fig2b.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0408572v1>

This figure "pietrzynski.fig3a.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0408572v1>

This figure "pietrzynski.fig3b.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0408572v1>